

LOGARITHMIC BOUNDS FOR ERGODIC SUMS OF CERTAIN FLOWS ON THE TORUS: A SHORT PROOF

JÉRÔME CARRAND

ABSTRACT. We give a short proof that the ergodic sums of \mathcal{C}^1 observables for a \mathcal{C}^1 flow on \mathbb{T}^2 admitting a closed transversal curve whose Poincaré map has constant type rotation number have growth deviating at most logarithmically from a linear one. For this, we relate the latter integral to the Birkhoff sum of a well-chosen observable on the circle and use the Denjoy-Koksma inequality. We also give an example of a nonminimal flow satisfying the above assumptions.

1. INTRODUCTION

Since the work of Furstenberg [12], it is known that the classical horocycle flow of a compact surface of constant negative curvature is uniquely ergodic — it has only one invariant Borel probability measure. This flow is related to a hyperbolic one, namely the geodesic flow, in the sense that the horocycle orbits are the unstable manifolds for the geodesic flow.

Using Symbolic Dynamics arguments (resp. equicontinuity of some functions), Marcus [16] (resp. [17]) generalized this result to the flow generated by the orientable one-dimensional unstable foliation of a connected basic piece of an Axiom A diffeomorphism (resp. flow). Later, Bowen and Marcus [4] extended this result to the higher dimensional strong stable or strong unstable foliation of a basic set for an Axiom A diffeomorphism or flow.

In their pioneer work, Giulietti and Liverani [13] focused on the one-dimensional stable foliation of a \mathcal{C}^r Anosov diffeomorphism F of the two-torus, inducing a flow h^t called the Giulietti–Liverani (stable horocycle) flow (of F). Giulietti and Liverani proved that this flow is uniquely ergodic, minimal and that it admits a closed transverse curve such that the rotation number of the first return map to this curve is of constant type. For more basic facts about this flow, see [2, Appendix A].

For any continuous function $f : \mathbb{T}^2 \rightarrow \mathbb{C}$, any $T > 0$ and any $x \in \mathbb{T}^2$, define the horocycle integral $H_{x,T}(f) = \int_0^T f(h^t(x)) dt$. By unique ergodicity, we have for any such x and f ,

$$\lim_{T \rightarrow \infty} \frac{H_{x,T}(f)}{T} = \mu^s(f) := \int_{\mathbb{T}^2} f d\mu^s,$$

where μ^s is the unique invariant probability measure of the flow h^t .

For large enough r , Giulietti and Liverani introduce a transfer operator for F on some suitable Banach space. Using eigenvectors of the dual operator associated

Date: August 17, 2022.

I thank S. Ghazouani for allowing me to use his idea for the proof and Y. Coudène for many useful comments. Research supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 787304).

to eigenvalues with modulus larger than the essential spectral radius (Ruelle resonances), they give an asymptotic expansion of $H_{x,T}(f)$ [13, Theorem 2.8]. The dominant term is the term $T\mu^s(f)$, corresponding to the trivial resonance $\lambda_0 = e^{h_{top}}$, where h_{top} is the topological entropy of F . This expansion also involves a negative power law error term. A simpler asymptotic expansion, in the case where all Ruelle resonances of the transfer operator have trivial Jordan blocks, can be found in [2, Equation (1.2)].

In their recent works, V. Baladi [2] and G. Forni [11] independently proved that horocycle integrals (in the set-up from [13]) do not have deviations, in other words the expansion is limited to the linear term with a bounded remainder. Their proofs are quite different: V. Baladi proves the strong result that the map F does not have non-trivial Ruelle resonance, while G. Forni uses the action of the (pseudo-)Anosov diffeomorphism on the first cohomology — in the more general setting of surfaces of genus $g \geq 1$ (non-trivial Ruelle resonances can appear only for $g \geq 2$).

In this note we give a new, much shorter, proof of the absence of deviations for horocycle integrals by considering a slightly more general setting: we no longer assume that the flow can be obtained from the stable foliation of an Anosov diffeomorphism. Instead, we only assume that the flow can be recovered from the suspension of a circle diffeomorphism whose rotation number is of constant type. In particular, these flows are uniquely ergodic. For clarity, we call “ergodic integral” for this type of flows the quantity defined as “horocycle integral” previously.

We give an elementary proof that the ergodic integral of a C^1 observable along the trajectory of such a flow on the two-torus grows at most logarithmically if the observable has zero average with respect to the unique invariant measure of the flow. This is the content of our main theorem (Theorem 2.2).

When comparing this estimate to the asymptotic expansion given by Giulietti and Liverani [13, Theorem 2.8], this result gives a new proof of the absence of deviations for the horocycle integral.

Finally, we prove that the class of flows we consider here is strictly larger than the class of flows studied by Giulietti and Liverani by constructing a flow satisfying our assumptions but which is not minimal — in contrast to all flows in [13]. This is the content of Theorem 3.1.

2. MAIN RESULT

Given a flow h_t on the two-torus, we call *ergodic integral* of an observable $f : \mathbb{T}^2 \rightarrow \mathbb{R}$ at $x \in \mathbb{T}^2$ and $T > 0$ the quantity $H_{x,T}(f) := \int_0^T f \circ h_t(x) dt$.

Recall the following classical theorem — we give a short proof of this fact using results from [15] in order to introduce notations for our main result. In particular the theorem below gives a simple sufficient condition for a flow to be written as the suspension of a circle diffeomorphism.

Theorem 2.1. *If h_t is a C^1 flow on the torus \mathbb{T}^2 without critical points nor periodic orbits, then there exists a smooth closed curve γ transverse to h_t such that h_t is smoothly conjugated to the suspension of the first return map $R : \gamma \rightarrow \gamma$. Moreover, the flow h_t is uniquely ergodic, with a unique invariant measure μ .*

Recall that an irrational number is of constant type if the sequence $(a_k)_k$ of its coefficients in its continued fraction expansion is bounded. We can now state our main result, using notations from the previous theorem.

Theorem 2.2. *If h_t is a C^1 flow on the torus \mathbb{T}^2 without critical point nor periodic orbit, and if the rotation number of the Poincaré first return map R is of constant type, then there exist constants K_1 and K_2 such that for any C^1 observable f with $\int f d\mu = 0$, any x and any $T > 0$,*

$$|H_{x,T}(f)| \leq K_1 \|f\|_{C^1} \log(1 + T) + K_2 \|f\|_{C^1}.$$

More precise versions of that estimate in the case of Giulietti–Liverani flows can be found in [2] and in [11]. The bound obtained by V.Baladi [2] is much tighter — but the proof is longer — while the estimate given by G.Forni [11] applies to flows on higher genus surfaces.

Proof of Theorem 2.1. By the Birkhoff recurrence theorem, any continuous transformation of a compact space has a recurrent point. Hence h_1 has recurrent orbits. In particular the flow h_t also has recurrent points. By our assumptions on the flow, these orbits cannot be periodic. Hence, by [15, Propositions 14.2.1 and 14.2.3] there exists a smooth closed curve γ transverse to h_t and parametrised by \mathbb{S}^1 such that every orbit of h_t intersects γ . We can therefore apply [15, Corollary 14.2.3] to get that h_t is smoothly conjugated to the suspension flow of the first return map R to γ . The conjugation is C^1 , since the change of coordinates is $(\theta, t) \mapsto h_t(\theta)$.

The map $R : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ is a C^1 diffeomorphism of the circle which has no periodic point. It is a classical result — see [6, Theorem 3.3.5] — that R is uniquely ergodic, with invariant measure ν , and that its rotation number is irrational. From this, we deduce that h_t is uniquely ergodic, with a unique invariant measure μ . \square

We can now give the proof of our main result.

Proof of Theorem 2.2. Suppose that the rotation number ω of R is of constant type. In order to prove the estimate, we will compare the ergodic integral to the Birkhoff sum of an appropriate function.

Let $u : \mathbb{S}^1 \rightarrow \mathbb{R}_+$ be the first return time function to γ , and let $f : \mathbb{T}^2 \rightarrow \mathbb{R}$ be a C^1 -observable such that $\int_{\mathbb{T}^2} f d\mu = 0$. By construction, γ is a smooth curve, uniformly transverse to the flow, hence the function u is of class C^1 . Define the C^1 observable g on γ by the formula

$$g(x) = \int_0^{u(x)} f \circ h_t(x) dt.$$

To estimate the ergodic integral of f by the Birkhoff sum of g under the map R , we use the following lemma.

Lemma 2.3. *For all $x \in \gamma$ and $T > 0$ there exists n satisfying $\frac{T}{\sup(u)} - 1 \leq n \leq \frac{T}{\inf(u)}$ and such that*

$$\left| H_{x,T}(f) - \sum_{k=0}^{n-1} g \circ R^k(x) \right| \leq \sup(u) \sup |f|.$$

For all $y \in \mathbb{T}^2$ there is $0 \leq \tau < \sup u$ and $x \in \gamma$ such that $y = h_\tau(x)$ and

$$|H_{x,T+\tau}(f) - H_{y,T}(f)| \leq \sup(u) \sup |f|.$$

Proof. We first determine n . Since $\inf u > 0$, there exists n such that $\sum_{k=0}^{n-1} u \circ R^k(x) \leq T < \sum_{k=0}^n u \circ R^k(x)$. Hence $n \inf u \leq T$ and $(n+1) \sup u \geq T$. Both estimates on ergodic integrals then follow from the fact that $h_t(R^n(x)) = h_{t+\sum_{k=0}^{n-1} u(R^k(x))}(x)$ for all $x \in \gamma$ and all $t \in \mathbb{R}$. \square

In order to conclude by applying the Denjoy–Koksma theorem [14, Theorem VI.3.1], we also need the following lemma.

Lemma 2.4. *If $\omega = [0, a_1, \dots, a_k, \dots]$ is of constant type, then for any integer $n > 1$ there exists integers N and (n_1, \dots, n_N) such that $n - 1 = \sum_{k=0}^N n_k q_k$, where $\frac{p_k}{q_k} = [0, a_1, \dots, a_k]$. Furthermore, we can choose $N < 4 \log(n) / \log(2)$ and $n_k \leq B$ for all k , where B is a bound on the coefficients $(a_k)_{k \geq 1}$.*

Proof. Since the sequence $(q_k)_{k \geq 0}$ satisfies the recursion formula $q_{k+1} = a_k q_k + q_{k-1}$ with $q_0 = 1$ and $q_1 = a_1$, we get by induction that $2^{\frac{k-1}{2}} \leq q_k$. Therefore, there exists N such that $q_N \leq n - 1 < q_{N+1}$ with the estimate $N < 4 \log(n) / \log(2)$.

Define inductively the sequences $(r_k)_{0 \leq k \leq N+1}$ and $(n_k)_{0 \leq k \leq N}$ by $r_{N+1} := n - 1$ and the Euclidean division $r_{k+1} = n_k q_k + r_k$, with $0 \leq r_k < q_k$. Clearly, we get that $n - 1 = \sum_{k=0}^N n_k q_k$ (because $q_0 = 1$). By contradiction, suppose there exists k such that $n_k > B + 1$. Then

$$r_{k+1} = n_k q_k + r_k > (B + 1)q_k + r_k > a_{k+1} q_k + q_{k-1} + r_k = q_{k+1} + r_k.$$

Therefore $r_{k+1} \geq q_{k+1}$, which is a contradiction. Hence $n_k \leq B$ for all k . \square

For completeness, we state the Denjoy–Koksma inequality:

Theorem 2.5 (Denjoy–Koksma inequality). *Let f be a homeomorphism of the circle with an irrational rotation number $\rho(f)$. Let μ be a measure invariant by f , and let p/q be such that $\gcd(p, q) = 1$ and $|q\rho(f) - p| < 1/q$. Then for all potential φ of bounded variation and all $x \in \mathbb{S}^1$, $\left| \sum_{k=0}^{q-1} \varphi \circ f^k(x) - q \int \varphi d\mu \right| < \text{Var}(\varphi)$.*

Since g is \mathcal{C}^1 , it is of bounded variation. In addition, the denominators $(q_k)_{k \geq 0}$ associated to ω satisfy the assumption $|q_k \omega - p_k| < 1/q_k$ for some integer p_k coprime with q_k . We can therefore apply the Denjoy–Koksma theorem to g , R and any q_k . Furthermore notice that, by construction, g is of ν -average 0: indeed, let $M = \{(x, t) \mid x \in \gamma, t \in [0, u(x)]\} / \sim$, with $(x, u(x)) \sim (R(x), 0)$, be the space such that h_t is conjugated with its unit speed vertical flow. Let $\bar{\mu}$ be the image of μ by the conjugacy map. Thus, $\bar{\mu}$ is invariant by the vertical flow and so it must be of the form $\bar{\mu} = \frac{1}{\int u d\nu} \bar{\nu} \otimes dt$, where $\bar{\nu}$ is invariant under R . By unique ergodicity of R , we have $\bar{\nu} = \nu$. Thus

$$\begin{aligned} 0 &= \int_{\mathbb{T}^2} f d\mu = \int_M f(h_t(x)) d\bar{\mu}(x, t) \\ &= \frac{1}{\int u d\nu} \int_{\gamma} \int_0^{u(x)} f(h_t(x)) dt d\nu(x) = \frac{1}{\int u d\nu} \int g d\nu. \end{aligned}$$

Fix $x \in \mathbb{T}^2$ and $T > 0$. By Lemma 2.3, there exist a point $y \in \gamma$ and an integer n from which we can estimate the ergodic integral of f at x and T with the Birkhoff sum of R at y . In order to assume that $n > 1$, we assume that $T > 2 \sup u$ (otherwise, the theorem holds with $K_1 = 0$ and some $K_2 > 0$ depending only on u). By Lemma 2.4 we can decompose $n - 1$ as a sum from which we deduce the equality

$$\sum_{k=0}^{n-1} g \circ R^k(y) = \sum_{l=0}^N \sum_{m=0}^{n_l-1} \sum_{k=0}^{q_l-1} g \circ R^k \left(R^{mq_l + \sum_{i=0}^{l-1} n_i q_i} y \right).$$

From the Denjoy-Koksma inequality, for all $0 \leq l \leq N$, all $0 \leq m < n_l$ and all y in γ ,

$$\left| \sum_{k=0}^{q_l-1} g \circ R^k \left(R^{mq_l + \sum_{i=0}^{l-1} n_i q_i} y \right) \right| < \text{Var}(g),$$

we deduce the estimate

$$\left| \sum_{k=0}^{n-1} g \circ R^k(y) \right| \leq NB \text{Var}(g) \leq \frac{4B \text{Var}(g)}{\log 2} \log n \leq \frac{4B \text{Var}(g)}{\log 2} \log \frac{T}{\inf(u)}.$$

Hence the result,

$$\begin{aligned} |H_{x,T}(f)| &\leq |H_{x,T}(f) - H_{y,T-\tau}(f)| + \left| H_{y,T-\tau}(f) - \sum_{k=0}^{n-1} g \circ R^k(y) \right| + \left| \sum_{k=0}^{n-1} g \circ R^k(y) \right|, \\ &\leq \frac{4B \text{Var}(g)}{\log 2} \log \frac{T}{\inf(u)} + 2 \sup(u) \sup |f| =: \tilde{K}_1 \log T + \tilde{K}_2. \end{aligned}$$

We can bound the total variation $\text{Var}(g)$ by the product of the length of γ with $\|g'\|_{\mathcal{C}^0(\gamma)}$. By the definition of g , we get

$$\|g'\|_{\mathcal{C}^0(\gamma)} \leq \|u'\|_{\mathcal{C}^0(\gamma)} \|f\|_{\mathcal{C}^0} + \|u\|_{\mathcal{C}^0(\gamma)} \|df\|_{\mathcal{C}^0} \sup_{0 \leq t \leq \|u\|_{\mathcal{C}^0(\gamma)}} \|dh_t\|_{\mathcal{C}^0}.$$

Notice that $\|u'\|_{\mathcal{C}^0(\gamma)}$ and $\sup_{0 \leq t \leq \|u\|_{\mathcal{C}^0(\gamma)}} \|dh_t\|_{\mathcal{C}^0}$ only depend on the flow h_t and on γ . Hence there exist constants K_1 and K_2 that depend only on h_t such that $\tilde{K}_1 \leq K_1 \|f\|_{\mathcal{C}^1}$ and $\tilde{K}_2 \leq K_2 \|f\|_{\mathcal{C}^1}$. \square

Finally, remark that in order to get a rotation number of constant type, the condition for the flow not to have periodic orbit is necessary: otherwise the existence of a transverse curve γ is no longer guaranteed. If such a curve exists then the first return map R has a periodic point, hence has a rational rotation number.

3. A NONMINIMAL FLOW SATISFYING THE ASSUMPTIONS OF THEOREM 2.2

We finish this note by proving that the class of flows we are working with is strictly larger than the class of flows studied by Giulietti and Liverani which are necessarily minimal. The proof relies on constructing a family of \mathcal{C}^1 nonminimal flows. By [15, Proposition 14.2.4], these flows are less than \mathcal{C}^2 .

Theorem 3.1. *There exists a flow on \mathbb{T}^2 satisfying the assumptions of Theorem 2.2 that is not minimal. Furthermore, the flow can be chosen to be renormalized by an Axiom A diffeomorphism.*

Notice however that all flows satisfying the assumptions of Theorem 2.2 are obtained by suspending circle diffeomorphisms of irrational rotation numbers, and thus are minimal on the support of their unique invariant measure.

Without the last condition of renormalization, we can simply construct such a flow by taking the suspension of a Denjoy counter-example whose rotation number is of constant type. Such circle diffeomorphisms exist by the original construction of Denjoy, which works for any irrational rotation number. For an expository on the construction of Denjoy counter-examples, see for example¹ [1]. However, there is no reason for the flow obtained by suspending a Denjoy counter-example to be renormalized by an Axiom A diffeomorphism. Adding this condition, the flow falls into the category of W^u -flows studied by Marcus in [16], in the particular case where the phase space of the flow is the same as the one of the Axiom A map — in opposition with just the set of nonwandering points of the map. Finally, results on Ruelle spectrum and dynamical determinants for Axiom A diffeomorphisms can be found in [3, 9] (and results on dynamical zeta functions for Axiom A flows in [10]), but asymptotic expansions of ergodic integrals associated to W^u -flows using transfer operator techniques are still quite rare in literature and there is room for work to be done in this setting.

In order to build a flow satisfying this last condition, consider the derived from Anosov transformation on the two-torus studied in [8, Chapter 9] and [7]. Recall some notation. Starting from Arnold's *cat map* (case $\beta = 0$) in the diagonalized form, and adding a bump in the unstable direction, let $f_\beta : [-\frac{1}{2}, \frac{1}{2}]^2 \rightarrow \mathbb{R}^2$ be as follows

$$f_\beta \begin{pmatrix} x \\ y \end{pmatrix} := \frac{1}{1+\lambda^2} \begin{pmatrix} \lambda & -1 \\ 1 & \lambda \end{pmatrix} \begin{pmatrix} \lambda^2 + \beta k \left(\frac{\sqrt{x^2+y^2}}{2} \right) & 0 \\ 0 & \lambda^{-2} \end{pmatrix} \begin{pmatrix} \lambda & 1 \\ -1 & \lambda \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix},$$

where $\lambda = \frac{1+\sqrt{5}}{2}$, $-\lambda^2 < \beta < 0$ and k is an even, unimodal function supported in $[-1, 1]$ such that $k(0) = 1$ — e.g. $k(r) = (1-r^2)^2 \mathbb{1}_{[-1,1]}(r)$ — so that the map f_β is invariant by the action of \mathbb{Z}^2 and induces a map, also called f_β , on the torus \mathbb{T}^2 . It is shown in [8, Chapter 9] that f_β is a diffeomorphism of class C^1 of the torus and if $-\lambda^2 < \beta < -\lambda^2 + 1$ then the origin is an attractive hyperbolic fixed point. Let K_β be the invariant subset defined as the complement of the basin of attraction of 0. This map is an explicit example of Smale's derived from Anosov transformation as introduced in [18, Section I.9], here obtained by perturbing Arnold's *cat map*.

Let $e_u = \frac{1}{\sqrt{1+\lambda^2}} \begin{pmatrix} \lambda \\ 1 \end{pmatrix}$ and $e_s = \frac{1}{\sqrt{1+\lambda^2}} \begin{pmatrix} -1 \\ \lambda \end{pmatrix}$ be unitary eigenvectors of the matrix $A := \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$ respectively associated to eigenvalues λ^2 and λ^{-2} . Since A is symmetric, notice that (e_u, e_s) is an orthonormal basis. In this basis the Jacobian matrix of f_β is

$$\text{Jac}(f_\beta)(x) = \begin{pmatrix} a_\beta(x) & b_\beta(x) \\ 0 & \lambda^{-2} \end{pmatrix}.$$

Since the Jacobian is upper-triangular, lines spanned by e_u are stable by f_β . Assuming that k satisfies also $k + id k' \leq 1$, $f_\beta|_{K_\beta}$ expands uniformly the direction

¹I thank Selim Ghazouani for indicating me this reference.

spanned by e_u . In order to construct a stable foliation over K_β , for X a vector field, denote $(f_\beta)_*X(x) = (d_x f_\beta)^{-1}X(f(x))$ to be the pullback of X by f_β . Formally, if $v_\beta^s = \lim_{n \rightarrow +\infty} \lambda^{-2n}(f_\beta)_*^n X$, then $\lambda^{-2}(f_\beta)_*v_\beta^s = v_\beta^s$, or in other words $d_x f_\beta v_\beta^s(x) = \lambda^{-2}v_\beta^s(f(x))$, v_β^s is uniformly contracted by $d f_\beta$. For the constant vector field $X \equiv e_s$, formally we get

$$(3.1) \quad v_\beta^s(x) = e_s - \sum_{i=0}^{\infty} \lambda^{-2i} b_\beta(f_\beta^i(x)) \prod_{j=0}^i \frac{1}{a_\beta(f_\beta^j(x))} e_u, \quad x \in \mathbb{T}^2.$$

This equation being only formal, we need to check that the series inside it converges. Since b_β is bounded and $a_\beta > 1$ on the compact set K_β , (3.1) defines a vector field on K_β , uniformly contracted by f_β :

$$(3.2) \quad d_x f_\beta v_\beta^s(x) = \lambda^{-2}v_\beta^s(f_\beta(x))$$

for all $x \in K_\beta$. It is shown in [5, Theorems 3.3 and 3.6] — in a slightly more general context — that (3.1) defines a Lipschitz continuous vector field on \mathbb{T}^2 for any fixed β in $] -\lambda^2 + \lambda^{-4}, 0]$ and that the map $(x, \beta) \mapsto v_\beta^s(x)$ is continuous on $\mathbb{T}^2 \times] -\lambda^2 + \lambda^{-4}, 0]$. Let h_t be the flow generated by $v_{\beta_0}^s$ for some fixed $-\lambda^2 + \lambda^{-4} < \beta_0 < -\lambda^2 + 1$. In fact, if we choose for the function k any \mathcal{C}^2 unimodal and even function supported in $[-1, 1]$, equal to 1 at 0 and satisfying $k + idk' \leq 1$, the induced vector fields v_β^s enjoys the same properties as before, but they are also \mathcal{C}^1 — see the discussion in [5, Theorem 3.7] — hence the flow h_t is also \mathcal{C}^1 . We make such a choice for k . We claim that this flow h_t satisfies the condition of Theorem 2.2 and that it is not minimal.

In order to prove this result, we first construct a closed transversal curve γ . We then construct a particular homotopy between the first return map and a rigid rotation, where none of the in-between map has a periodic point. From the continuity of the rotation number, it is enough to compute the rotation number of the rigid rotation, which happens to be a quadratic integer. The nonminimality follows from the invariance of the proper closed set K_{β_0} by the flow h_t . First we need the following lemma.

Lemma 3.2. *The flow h_t does not have periodic orbit. This is also true for the flow generated by v_β^s for any $-\lambda^2 + \lambda^{-4} < \beta \leq 0$.*

Proof. By construction, each vector field v_β^s satisfies $d_x f_\beta(v_\beta^s(x)) = \lambda^{-2}v_\beta^s(f_\beta(x))$. By differentiating $f_{\beta_0} \circ h_t(x)$ and $h_{\lambda^{-2}t} \circ f_{\beta_0}(x)$ according to t , we get that these two functions satisfy the same Cauchy problem for all $x \in \mathbb{T}^2$, thus the relation

$$(3.3) \quad f_{\beta_0} \circ h_t = h_{\lambda^{-2}t} \circ f_{\beta_0}$$

holds by uniqueness of the solution (because v_β^s is Lipschitz continuous). Therefore, if by contradiction h_t has a periodic orbit, by applying $f_{\beta_0}^n$, for n large enough, we get an arbitrarily short periodic orbit for the flow. This contradicts the fact that the component along e_s in the basis (e_u, e_s) of $v_{\beta_0}^s$ is constant equal to 1. \square

Proof of Theorem 3.1. Since the map $(x, \beta) \mapsto v_\beta^s(x)$ is continuous on the compact set $\mathbb{T}^2 \times [\beta_0, 0]$, the component of these vector fields in the basis (e_u, e_s) along e_u is uniformly bounded and along e_s is equal to 1, by definition. Therefore, there exists a vector w of rational slope, say $w = \frac{1}{\sqrt{p^2+q^2}} \begin{pmatrix} q \\ p \end{pmatrix}$, where p and q are coprimes, so

that w is uniformly transverse to v_β^s for all $\beta \in [\beta_0, 0]$. Define γ to be the closed curve passing through $(0, 0)$ and with slope p/q . By choice of w , the curve γ is transverse to v_β^s and so for every β in $[\beta_0, 0]$. We can naturally parametrize γ by \mathbb{S}^1 .

Let $R : \mathbb{S}^1 \rightarrow \mathbb{S}^1$ be the first return map to γ of h_t . Notice that performing a time change on this flow does not affect the first return map R , but only the first return time function u . In order to simplify computations, renormalize the vector fields as follows

$$w_\beta^s = \frac{1}{\langle v_\beta^s, w^\perp \rangle} v_\beta^s$$

so that, for each β , the flow $\phi_t^{(\beta)}$ generated by w_β^s has a constant first return time function $u_\beta \equiv \tau_\beta$, where w^\perp is the unitary vector equal to w rotated by an angle $\pi/2$. These first return time functions do not depend on β , in other words $\tau_\beta \equiv \tau$. Since $b_0 \equiv 0$, notice that w_0^s is a constant vector field (equals everywhere to e_s), hence its first return map to γ is a rigid translation $R_\alpha : x \mapsto x + \alpha$. Introduce also the notation $R^{(\beta)}$ for the first return map to γ of $\phi_t^{(\beta)}$. In particular $R = R^{(\beta_0)}$ and $R_\alpha = R^{(0)}$.

By [5, Theorem 3.10], the map $\beta \mapsto v_\beta^s$ is continuous for the \mathcal{C}^0 -topology on the space of vector fields. From a Gronwall type argument, we get that $\beta \mapsto R^{(\beta)}$ is continuous for the \mathcal{C}^0 -topology. Now, by [14, Proposition II.2.7], the map $\beta \mapsto \rho(R^{(\beta)})$ is continuous, where $\rho(R^{(\beta)})$ stands for the rotation number of $R^{(\beta)}$. In order to prove that $\rho(R) = \alpha$, we prove that $\rho(R^{(\beta)})$ cannot be rational, but this directly follows from Lemma 3.2. Hence $\beta \mapsto \rho(R^{(\beta)})$ is a constant map and $\rho(R) = \alpha$.

We now compute the value of α . Consider lifts $\tilde{w}_{(0)}^s$, $\tilde{\gamma}$ and $\tilde{\phi}_t^{(0)}$ to \mathbb{R}^2 of respectively w_0^s , γ and $\phi_t^{(0)}$. Let (∂_x, ∂_y) be the canonical basis of \mathbb{R}^2 . Notice that the arc $\{\tilde{\phi}_t^{(0)}((0, 1)) \mid -p\tau \leq t \leq 0\}$ starts at the point $(0, 1)$ and ends on the branch of $\tilde{\gamma}$ containing $(0, 0)$ at some point cw , for some $c > 0$. The coordinates of this intersection point satisfy the system of equations

$$\begin{cases} -p\tau \langle w_{(0)}^s, \partial_x \rangle = cq(p^2 + q^2)^{-1/2} \\ 1 - p\tau \langle w_{(0)}^s, \partial_y \rangle = cp(p^2 + q^2)^{-1/2}, \end{cases}$$

where $\langle \cdot, \cdot \rangle$ denotes the usual scalar product. Now, notice that $c/|\gamma| = -p\alpha$, where $|\gamma|$ is the length of the closed curve γ . We can solve these equations for α and get

$$\alpha = \frac{1}{pq} \frac{1}{\lambda - \frac{p}{q}}$$

which clearly is a quadratic integer, since λ is. Therefore α is of constant type.

The nonminimality of h_t is ensured by properties proven in [8, Chapter 9]. More precisely, let U be the basin of attraction of $(0, 0)$ for f_{β_0} and K be its complement in the torus. In [8, Chapter 9], Coudène proved that the set K is nonempty and that U and K are invariant by f_{β_0} . Now, because of (3.3), the sets U and K are invariant by the flow h_t .

Finally, the map f is an Axiom A diffeomorphism since f is transitive [8, Chapter 9] on the hyperbolic set K [5, Theorem 2.9]. Therefore, by the shadowing lemma,

periodic points are dense in the compact invariant set K which coincides with the nonwandering set of f . \square




FIGURE 1. Representation of the minimal component K of the flow (h_t) . Underneath is the vector field v^s generating the flow.

Finally, we give in Figure 1 a representation of the set K . In [8, Chapter 9], it is proven that K is the closure of the stable leaf $W^s(p)$ of a hyperbolic fixed point p for f_{β_0} . From the relation (3.3) and the Hartman-Grobman theorem, it follows that this stable leaf is equal to the orbit of p by the flow h_t . From [6, Theorem 3.3.4], the set $K \cap \gamma$ coincides with any ω -limit set and any α -limit set of R . Therefore, the set K is the minimal component of h_t and is also an attractor for both positive and negative times. Moreover, K is also the support of the unique invariant measure μ of h_t .

APPENDIX A. ALTERNATIVE PROOF OF THEOREM 3.1 FROM SEMI-CONJUGACY

We give an alternative proof of Theorem 3.1. More precisely, we use the same example, but we compute the rotation number in a different way: we construct a semi-conjugacy map h so that $h \circ R = R_\alpha \circ h$. It will follow that the rotation number of R is α . The construction of h is inspired from the proof of [19, Proposition 7].

Proof. Exactly as in the first proof of Theorem 3.1, we construct the closed transversal curve γ and we renormalize the vector fields v_β^s so that the time of first return function to γ of their associated flows is constant. The computation of α remains the same, and we get that α is a quadratic integer, hence α is of constant type. In particular, the rotation R_α is minimal.

We now prove that the first return map R of h_t is semi-conjugated to R_α . To this end, we construct a surjective and continuous function h of the circle.

Let $h(R^n(0)) := R_\alpha^n(0)$ for all $n \in \mathbb{Z}$. This map is well defined since h_t has no periodic orbit by Lemma 3.2, so does R . In order to extend h into a continuous map, we first prove that it preserves order of triplets. Fix an orientation of \mathbb{S}^1 — and therefore of γ — seen as \mathbb{R}/\mathbb{Z} . Let $x_1 := R^{n_1}(0)$, $x_2 := R^{n_2}(0)$ and $x_3 := R^{n_3}(0)$

be so that (x_1, x_2, x_3) is an ordered triplet of \mathbb{S}^1 — we can assume that n_1, n_2 and n_3 are distinct. We prove that the triplet $(x'_1, x'_2, x'_3) = (h(x_1), h(x_2), h(x_3))$ is also ordered. Consider the family of curves $\varphi_\beta := \{\phi_t^{(\beta)}(0) \mid \min(n_1, n_2, n_3)\tau \leq t \leq \max(n_1, n_2, n_3)\tau\}$. By continuity of $(x, \beta) \mapsto w_\beta^s(x)$, this family depends on β in a continuous fashion.

Notice that points x_1, x_2 and x_3 correspond to some intersection points between φ_{β_0} and γ , and that points x'_1, x'_2 , and x'_3 correspond to some intersection points between φ_0 and γ . Furthermore, we can connect x_1 to x'_1 (respectively x_2 to x'_2 , and x_3 to x'_3) with intersection points between γ and φ_β when varying the value of β . Therefore we can track the evolution of (x_1, x_2, x_3) with continuous functions $(x_1(\beta), x_2(\beta), x_3(\beta))$ of β such that $x_1(\beta_0) = x_1$ and $x_1(0) = x'_1$ — and similarly for $x_2(\beta)$ and $x_3(\beta)$.

By contradiction, suppose that the triplet (x'_1, x'_2, x'_3) is not ordered. By continuity, this means that for some value of β_1 in $[\beta_0, 0]$ and without loss of generality $x_1(\beta_1) = x_2(\beta_1)$. In other words, this means that the first return map to γ of $\phi_t^{(\beta_1)}$ has a periodic point, which contradicts Lemma 3.2.

Therefore, the map h can be lifted into a “degree” one, increasing, function $\tilde{h} : \pi^{-1}\{R^n(0) \mid n \in \mathbb{Z}\} \rightarrow \pi^{-1}\{R_\alpha^n(0) \mid n \in \mathbb{Z}\}$, where $\pi : \mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z}$ is the canonical projection. In other words, $\pi \circ \tilde{h} = h \circ \pi$ and $\tilde{h}(x+1) - \tilde{h}(x) = 1$ for all x where \tilde{h} is defined. By minimality of R_α , the range of \tilde{h} is dense in \mathbb{R} . Hence, we can uniquely extend \tilde{h} by a continuous, increasing and surjective function $\tilde{h} : \mathbb{R} \rightarrow \mathbb{R}$. Its projection on the circle, still noted h , is also continuous and extends h into a degree one map of the circle. By continuity of R and of R_α , we get that $h \circ R = R_\alpha \circ h$. Therefore, by [14, Proposition II.2.10], the rotation number of R is α , a quadratic integer.

The nonminimality of h_t is ensured by properties proven in [8, Chapter 9]. \square

Remark A.1. The construction of the conjugacy map h comes from the following heuristic. Since the stable manifold of 0 under the cat map is blown up into an open set, the basin of attraction $U_\beta := \mathbb{T}^2 \setminus K_\beta$ of 0 under f_β , we expect that the map h relates the orbit of 0 under R_α with the orbit of I under R , where I is the connected component of $\gamma \cap U_\beta$ containing 0 (notice that I is a wandering interval and that its orbit under R is $\gamma \cap U_\beta$, which is dense in γ). More precisely, we expect h to be similar to the Cantor staircase function, being constant when restricted to each $R^n(I)$. As in the construction of the Cantor staircase function, we only need to know the values of h where it is constant, as long as h is non-decreasing and that this set of values has a connected closure. In the proof above, we chose to define h first by setting $h(x_n) = R_\alpha^n(0)$ with $x_n = R^n(0)$, but we could have chosen any sequence $x_n \in R^n(I)$.

REFERENCES

- [1] K. Athanassopoulos. Denjoy C^1 diffeomorphisms of the circle and McDuff’s question. *Expositiones Mathematicae*, 33(1):48–66, 2015.
- [2] V. Baladi. There are no deviations for the ergodic averages of Giulietti–Liverani horocycle flows on the two-torus. *Ergodic Theory and Dynamical Systems*, pages 1–14, 2019.
- [3] V. Baladi and M. Tsujii. Dynamical determinants and spectrum for hyperbolic diffeomorphisms. In *Geometric and probabilistic structures in dynamics*, volume 469 of *Contemp. Math.*, pages 29–68. Amer. Math. Soc., Providence, RI, 2008.
- [4] R. Bowen and B. Marcus. Unique ergodicity for horocycle foliations. *Israel J. Math.*, 26(1):43–67, 1977.

- [5] J. Carrand. Explicit construction of non-linear pseudo-Anosov maps, with nonminimal invariant foliations. *arXiv preprint arXiv:2104.11625*, 2021.
- [6] I. P. Cornfeld, S. V. Fomin, and Y. G. Sinai. *Ergodic Theory*, volume 245. Springer Science & Business Media, 2012.
- [7] Y. Coudène. Pictures of hyperbolic dynamical systems. *Notices of the AMS*, 53(1), 2006.
- [8] Y. Coudène. *Ergodic Theory and Dynamical Systems*. Springer, 2016.
- [9] N. V. Dang and G. Rivière. Pollicott-Ruelle spectrum and Witten Laplacians. *J. Eur. Math. Soc. (JEMS)*, 23(6):1797–1857, 2021.
- [10] S. Dyatlov and C. Guillarmou. Afterword: dynamical zeta functions for Axiom A flows. *Bull. Amer. Math. Soc. (N.S.)*, 55(3):337–342, 2018.
- [11] G. Forni. On the equidistribution of unstable curves for pseudo-Anosov diffeomorphisms of compact surfaces. *arXiv preprint arXiv:2007.03144*, 2020.
- [12] H. Furstenberg. The unique ergodicity of the horocycle flow. In *Recent advances in topological dynamics (Proc. Conf., Yale Univ., New Haven, Conn., 1972; in honor of Gustav Arnold Hedlund)*, pages 95–115. Lecture Notes in Math., Vol. 318, 1973.
- [13] P. Giulietti and C. Liverani. Parabolic dynamics and Anisotropic Banach spaces. *Journal of the European Mathematical Society*, 21(9):2793–2858, 2019.
- [14] M. R. Herman. Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations. *Publications Mathématiques de l’IHÉS*, 49:5–233, 1979.
- [15] A. Katok and B. Hasselblatt. *Introduction to the modern theory of dynamical systems*, volume 54 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1995. With a supplementary chapter by Katok and Leonardo Mendoza.
- [16] B. Marcus. Unique ergodicity of some flows related to Axiom A diffeomorphisms. *Israel J. Math.*, 21(2-3):111–132, 1975.
- [17] B. Marcus. Unique ergodicity of the horocycle flow: variable negative curvature case. *Israel J. Math.*, 21(2-3):133–144, 1975.
- [18] S. Smale. Differentiable dynamical systems. *Bulletin of the American Mathematical Society*, 73(6):747–817, 1967.
- [19] J.-C. Yoccoz. Echanges d’intervalles. *Cours Collège de France*, https://www.college-de-france.fr/media/jean-christophe-yoccoz/UPL8726_yoccoz05.pdf, 2005.

LABORATOIRE DE PROBABILITÉS, STATISTIQUES ET MODÉLISATION (LPSM), CNRS, SORBONNE UNIVERSITÉ, UNIVERSITÉ DE PARIS, 4, PLACE JUSSIEU, 75005 PARIS, FRANCE

Email address: jcarrand@lpsm.paris